

兔抗 BCAM 多克隆抗体

中文名称：兔抗 BCAM 多克隆抗体

英文名称： Anti-BCAM rabbit polyclonal antibody

别 名： AU; LU; CD239; MSK19

相关类别：一抗

抗 原：BCAM

储 存：冷冻 (-20°C)

宿 主：Rabbit

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

This gene encodes Lutheran blood group glycoprotein, a member of the immunoglobulin superfamily and a receptor for the extracellular matrix protein, laminin. The protein contains five extracellular immunoglobulin domains, a single transmembrane domain, and a short C-terminal cytoplasmic tail. This protein may play a role in epithelial cell cancer and in vaso-occlusion of red blood cells in sickle cell disease. Polymorphisms in this gene define some of the antigens in the Lutheran system and also the Auberger system. Inactivating variants of this gene result in the recessive Lutheran null phenotype, Lu(a-b-), of the Lutheran blood group. Tw

	o transcript variants encoding different isoforms have been found for this gene.
Applications:	ELISA, WB
Name of antibody:	BCAM
Immunogen:	Fusion protein of human BCAM
Full name:	basal cell adhesion molecule (Lutheran blood group)
Synonyms :	AU; LU; CD239; MSK19
SwissProt:	P50895
ELISA Recommended dilution:	1000-2000
WB Predicted band size:	67 kDa
WB Positive control:	293T and 231 cells
WB Recommended dilution:	200-1000

